WOMEN & ALCOHOL

Free Full section on Women and Alcohol Use Disorder (AUD), taken from the book.

The complete book includes dedicated material about women and alcohol alongside behavioural, psychological and medical approaches to change.



Alcohol: Cut Down. Stop. Stay Stopped

by J.S McInally

APPENDIX G: WOMEN & AUD

Biology, psychology, pressure and the double standard

If you're a woman and alcohol has started to take more from you than it gives, you're not imagining it, and you're not "weak". There are real, measurable reasons alcohol can hit women harder, progress faster and feel harder to quit, even when the drinking looks "less severe" on the outside.

And then there's the second layer: the social judgement. Women are often judged more harshly for the same behaviour, expected to cope quietly, expected to stay functional, expected to keep everyone else steady. When alcohol becomes the relief valve, it can feel like the only private place you get to exhale.

This chapter is here to explain what's true, what's evidence-based and what's changeable. Not to frighten you. Not to shame you. To give you clarity — and then a way forward.

Why alcohol can affect women more than men

Let's start with the body, because this part matters and it's grounded in biology.

Higher blood alcohol levels, faster metabolisation

On average, women have a higher proportion of body fat and less total body water than men. Alcohol disperses mainly in body water. This means

that, pound for pound, the same amount of alcohol often leads to a higher blood alcohol concentration in women. The result is stronger effects, earlier impairment and greater physiological strain, even when drinking the same number of units.

Alcohol-related harm can appear earlier

Research consistently shows that some women develop alcohol-related problems after a shorter drinking history and at lower levels of consumption compared to men. This pattern is sometimes described clinically as a faster progression of harm. It does not affect all women in the same way, but as a population trend it is well recognised in medicine and addiction science.

Cancer risk is not theoretical

Alcohol increases the risk of several cancers, including breast cancer. This matters because alcohol can influence hormone levels such as oestrogen and can contribute to biological processes involved in tumour development. This risk exists regardless of whether the alcohol comes from wine, spirits or beer.

These differences don't mean women are fragile. They mean women are biologically different — and alcohol does not affect all bodies equally.

Why so many women drink: the emotional logic behind it

People don't keep doing something that harms them unless it also gives them something they feel they need.

For many women, alcohol begins as a solution before it becomes a problem. It offers short-term relief from very real pressures: anxiety, insomnia, emotional overload, loneliness and burnout. In that sense, drinking often makes psychological sense long before it becomes destructive.

Common reasons women report drinking include:

- switching off anxiety and constant mental noise

- getting to sleep
- numbing emotional pain or loneliness
- coping with relentless responsibility
- rewarding themselves after holding everything together
- managing shame, self-criticism or exhaustion
- tolerating situations or relationships that feel emotionally unsafe

None of these reasons reflect weakness. They reflect adaptation.

Alcohol temporarily activates calming and reward pathways in the brain. But that relief is followed by rebound anxiety, disrupted sleep and low mood. Over time, the brain begins to rely on alcohol not just for pleasure, but for emotional regulation. The very symptoms alcohol was used to treat then become the triggers to drink again.

This is how the loop forms. And once it's established, willpower alone rarely breaks it.

Stigma, secrecy and why women often delay asking for help

Women tend to face a harsher moral judgement around alcohol problems than men. That judgement may be explicit — labels, gossip, criticism — or subtle: distance, raised eyebrows, quiet withdrawal of support.

The effect is the same. Stigma drives drinking underground. It encourages secrecy. It delays help-seeking. And it increases the likelihood of trying to "handle it privately", even when that becomes unsafe.

Many women describe intense fear of being judged as a failure — as a mother, partner, professional or caregiver. Others fear the consequences of disclosure: losing trust, status or stability. These fears, whether real or anticipated, keep many women silent far longer than is healthy.

This is not a lack of insight. It is the predictable result of stigma combined with responsibility.

Sexism, gender pressure and how they feed drinking

Sexism does not need to be overt to shape behaviour. Often it works quietly, through expectations that are impossible to meet.

Many women live under constant pressure to be:

- capable but not intimidating
- attractive but not self-absorbed
- emotionally available but not "needy"
- resilient but not demanding
- productive while carrying the emotional load for others

This ongoing self-monitoring is exhausting. Alcohol can feel like the fastest way to stop performing.

There is also the issue of emotional labour. Many women carry invisible work: managing relationships, anticipating needs, smoothing conflict, holding families together and keeping life running. When the day finally ends, the nervous system may still be fully switched on. Alcohol becomes the off switch.

Trauma also plays a role for many women. Rates of certain forms of interpersonal trauma are higher among women, and alcohol is a common way of dampening a nervous system stuck in threat mode. This is not about blame. It is about recognising how survival strategies become habits.

Add to this a culture that markets alcohol to women as self-care — "wine o'clock", "mum needs a drink", "you deserve it" — and it becomes clear that many women are not drinking irresponsibly. They are responding to pressure with the tools they've been given.

Hormones, life stages and vulnerability

Hormones do not cause alcohol use disorder on their own. But they can strongly influence mood, sleep, anxiety and stress sensitivity — all of which affect drinking behaviour.

The menstrual cycle

Fluctuations in oestrogen and progesterone across the cycle can affect emotional regulation, impulse control, sleep quality and anxiety levels. Some women notice stronger cravings or reduced tolerance during certain phases of the month. Understanding these patterns can make urges feel less mysterious and more manageable.

Perimenopause and menopause

Perimenopause can bring disrupted sleep, anxiety, irritability, low mood, cognitive fog and a sense of not feeling like yourself. Alcohol may appear to help initially, especially with sleep and tension.

Over time, however, alcohol often worsens sleep quality and increases next-day anxiety. For some women, it also intensifies hot flushes and night sweats. Midlife can also coincide with major stressors: caring responsibilities, relationship strain, identity shifts, grief and career pressure. If alcohol has been a coping tool for years, these converging factors can dramatically increase reliance on it.

The goal during this stage is not perfection. It is stability. Supporting sleep, nervous system regulation and emotional containment become essential.

Pregnancy

There is no known safe level of alcohol during pregnancy. Alcohol exposure can lead to fetal alcohol spectrum disorder, which is entirely preventable. For women who are already dependent, pregnancy requires medical support.

Abruptly stopping alcohol can be dangerous, and safety must always come first.

Why women can look “fine” while suffering deeply

Many women continue to function at a high level while struggling intensely.

- They work.
- They care for others.
- They keep up appearances.
- And then they collapse privately.

Functioning does not equal wellness. It often means someone has become very good at hiding distress — including from themselves. Alcohol use disorder is not defined by external chaos. It is defined by the relationship with alcohol, the loss of control, the psychological cost and the internal toll.

If drinking is secretive, compulsive, planned around, followed by shame or anxiety, or used to manage emotions — it matters. Even if everything else still looks intact.

What supports recovery for women

Women do not need a simplified or softened version of recovery. They need approaches that reflect real risk factors and real barriers.

Key elements that consistently support change include:

Safety first: Alcohol withdrawal can be dangerous. Anyone with physical dependence needs a planned, supported approach.

Reducing shame: Shame delays recovery. Honest, non-judgemental support accelerates it.

Addressing sleep and anxiety early: These are not side issues. For many women they are the core drivers of drinking.

Practical support that fits real lives: Work, caregiving and financial pressure are not excuses. They are structural realities that recovery plans must accommodate.

Understanding the 'why', without waiting for it: Deeper emotional work matters, but early recovery begins with stabilising the body and brain. Feeling better first makes change possible.

Putting This Into Context

Many women carry a quiet belief that they should be able to cope. That if they're struggling, it means they've failed. The evidence tells a different story.

You are a nervous system under pressure. Alcohol reduced that pressure — temporarily. Your brain learned the pattern. Now it's stuck. The loop is not your fault. But it *is* breakable. Not through toughness. Through clarity, safety, structure, support and time. And if judgement or stigma has kept you silent, let this be the line in the sand: You do not have to earn help by suffering more.

You are allowed to get better now.